

MULTIMEDIA



UNIVERSITY

STUDENT ID NO

--	--	--	--	--	--	--	--	--	--

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 2, 2018/2019

DEM5028 – ENGINEERING MATHEMATICS 2

(Diploma in Electronic Engineering)

4 MARCH 2019

2.30 PM – 4.30 PM

(2 Hours)

INSTRUCTIONS TO STUDENT

1. This question paper consists of 3 pages (excluding the cover page and appendices).
2. Attempt **ALL FOUR** questions.
3. Write your answers in the answer booklet provided.
4. Formulae are provided in the appendix section.

QUESTION 1

a. Evaluate the following integrals.

i. $\int \frac{5x^2 + 20x + 6}{x^3 + 2x^2 + x} dx$ (8 marks)

ii. $\int_0^{\frac{\pi}{4}} x \sec^2 x \, dx$ (5 marks)

b. The Diagram 1 below shows a straight line $2x + y = 16$ intersecting a curve $y = (x - 4)^2$ at point P and point Q . Find




Diagram 1

- i. the area of the region A. (8 marks)
- ii. the volume generated when the shaded region B is revolved through 360° about the x-axis. (4 marks)

[TOTAL 25 MARKS]

Continued ...

QUESTION 2

a. Find the general solution of the differential equation $2 + \frac{dy}{dx} = \frac{3x}{y-2} + 2$. (5 marks)

b. Find the solution of the differential equation $y^2 \frac{dy}{dx} = -3y\sqrt{x}$, given that $y = 1$ when $x = 1$. (9 marks)

c. Find the solution of differential equation $x \frac{dy}{dx} - x^4 = -y$, subject to $y(1) = 1$. (11 marks)

[TOTAL 25 MARKS]

QUESTION 3

a. Find the sum to infinity for the series (4 marks)

$$18 - 6 + 2 - \dots$$

b. Expand $(9 + x)^{\frac{1}{2}}$ in ascending powers of x up to the term in x^2 . (7 marks)

c. Given that $A(3, -1, 1)$, $B(-1, 2, -2)$ and $C(3, 2, -2)$ are three vertices of a triangle

- Find $\overrightarrow{AB} \times \overrightarrow{BC}$. (7 marks)
- Calculate the area of triangle ABC . (7 marks)

[TOTAL 25 MARKS]

Continued ...

QUESTION 4

a. If $f(x, y) = e^{2x} \sin(3x + 2y)$.

i. Compute $\frac{\partial f}{\partial x} \Big|_{\left(0, \frac{1}{8}\pi\right)}$ and $\frac{\partial f}{\partial y} \Big|_{\left(0, \frac{1}{8}\pi\right)}$. (9 marks)

ii. Evaluate f_{xx} and f_{yy} . (8 marks)

b. Find the volume of the solid bounded above by the plane $z = 4 - x - y$ and below by the rectangle $R = \{(x, y) : 0 \leq x \leq 1, 0 \leq y \leq 2\}$. (8 marks)

[TOTAL 25 MARKS]

End of Page.

APPENDICES: Formulae**Integration of common functions**

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad \int \frac{1}{x} dx = \ln|x| + C \quad \int e^x dx = e^x + C \quad \int a^x dx = \frac{a^x}{\ln a} + C$$

$$\int \sin x dx = -\cos x + C \quad \int \cos x dx = \sin x + C \quad \int \sec^2 x dx = \tan x + C \quad \int \sec x \tan x dx = \sec x + C$$

$$\int \csc^2 x dx = -\cot x + C \quad \int \csc x \cot x dx = -\csc x + C$$

Inverse Trigonometry**Pythagorean Identities****Integration by parts**

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C$$

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \cot^2 x = \csc^2 x$$

$$\int u dv = uv - \int v du$$

$$\int \frac{1}{x^2+1} dx = \tan^{-1} x + C$$

$$1 + \tan^2 x = \sec^2 x$$

Areas Between Curves**Volume by Washer****Volume by Cylindrical Shells**

$$A = \int_a^b [f(x) - g(x)] dx$$

$$V = \int_a^b \pi([f(x)]^2 - [g(x)]^2) dx$$

$$V = \int_a^b 2\pi x(f(x) - g(x)) dx$$

$$A = \int_c^d [w(y) - v(y)] dy$$

$$V = \int_c^d \pi([w(y)]^2 - [v(y)]^2) dy$$

$$V = \int_c^d 2\pi y(w(y) - v(y)) dy$$

Linear Differential Equations:

$$\frac{dy}{dx} + p(x)y = q(x); \mu y = \int \mu q(x) dx \Rightarrow y = \frac{1}{\mu} \int \mu q(x) dx, \quad \text{where } \mu = e^{\int p(x) dx}$$

Divergence Test	If $\lim_{n \rightarrow \infty} a_n \neq 0$, then $\sum a_n$ diverges.
p-series	The p-series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ is convergent if $p > 1$ and divergent if $p \leq 1$.
Limit Comparison Test	Let $\sum a_n$ and $\sum b_n$ be series with positive terms such that $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$ If $0 < c < \infty$, then both series converge or both diverge.
Alternating Series Test	If the alternating series $\sum_{n=1}^{\infty} (-1)^{n-1} b_n = b_1 - b_2 + b_3 - b_4 + \dots \quad b_n > 0$ Satisfies : i. $b_{n+1} \leq b_n$ for all n ii. $\lim_{n \rightarrow \infty} b_n = 0$ then the series is convergent
Ratio Test	Let $\sum a_n$ be a series with nonzero terms such that $L = \lim_{n \rightarrow \infty} \frac{ a_{n+1} }{ a_n }$ a. Series converges absolutely if $L < 1$ b. Series diverges if $L > 1$ or $L = \infty$ c. No conclusion if $L = 1$

Binomial expansion

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

Vector

The length of the vector $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$ is $|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$.

If θ is the angle between the vector \mathbf{a} and \mathbf{b} , then $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}|\cos\theta$ & $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}||\mathbf{b}|\sin\theta$

Cross Product

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \mathbf{k}$$

Equation of Line

Vector equation: $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$

Parametric equation: $x = x_0 + at$, $y = y_0 + bt$, $z = z_0 + ct$

Equation of Plane $\langle a, b, c \rangle \cdot \langle x - x_0, y - y_0, z - z_0 \rangle = 0$

The Chain Rule

Suppose that $z = f(x, y)$, where $x = g(t)$ and $y = h(t) \Rightarrow \frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$

Second Derivatives Test

Suppose that $f_x(a, b) = 0$ and $f_y(a, b) = 0$ [that is, (a, b) is a critical point of f]. Let

$$D = D(a, b) = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2$$

- If $D > 0$ and $f_{xx}(a, b) > 0$, then $f(a, b)$ is a local minimum.
- If $D > 0$ and $f_{xx}(a, b) < 0$, then $f(a, b)$ is a local maximum
- If $D < 0$, then $f(a, b)$ is a saddle point.

Moments and Centers of Mass

The **moment** about the **x-axis**:

$$M_x = \iint_D y\rho(x, y)dA$$

The **moment** about the **y-axis**:

$$M_y = \iint_D x\rho(x, y)dA$$

The coordinates (\bar{x}, \bar{y}) of the center of mass:

$$\bar{x} = \frac{M_y}{m} = \frac{1}{m} \iint_D x\rho(x, y)dA \quad \bar{y} = \frac{M_x}{m} = \frac{1}{m} \iint_D y\rho(x, y)dA$$

Where the mass:

$$m = \iint_D \rho(x, y)dA$$

Triple Integrals: $\iiint_B f(x, y, z)dV = \int_r^s \int_c^d \int_a^b f(x, y, z)dx dy dz$